



Lipid hydrolysis products affect the composition of microbiota isolated from infant fecal samples after in vitro fermentation

Bennike, Rikke Mette Guldhammer; Licht, Tine Rask; Hellgren, Lars

Publication date:
2013

Document Version
Early version, also known as pre-print

[Link back to DTU Orbit](#)

Citation (APA):
Bennike, R. M. G., Licht, T. R., & Hellgren, L. (2013). *Lipid hydrolysis products affect the composition of microbiota isolated from infant fecal samples after in vitro fermentation*. Abstract from Cell Symposia, Lisbon, Portugal.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Abstract for Cell Symposia – Microbiome and Host Health
May 12-14 2013, Lisbon, Portugal

Lipid hydrolysis products affect the composition of microbiota isolated from infant fecal samples after *in vitro* fermentation.

Rikke MG Bennike, Tine R Licht, Lars I Hellgren

Some lipid hydrolysis products such as medium-chained free fatty acids (FFA) and monoacylglycerols (MAG) have antibacterial activity, while others, including oleic acid, have been reported to be essential for optimal growth of *Lactobacillus* species. Thus, the FFA and MAG concentration in the distal ileum and in colon can be expected to selectively modulate the growth rate and hereby the composition of the microbiota. In earlier studies, we have shown that this concentration is dependent on the type of emulsification of the triglycerides, which deviates between breast milk and formula milk.

Here, we have determined effects of selected combinations of FFA and MAG on microbial composition during a 24-hour anaerobic *in vitro* fermentation in microbiota obtained from infant fecal samples (age 2-5 months). PCR-based quantification of 11 different bacterial taxa revealed that the growth of *Firmicutes*, *Lactobacillus* and *B.longum* is significantly increased in the presence of a mixture of C10-C14 FFAs.